

## **Product Datasheet**

## HER2 Antibody / ErbB2 (orb2643198)

**Description** Recognizes a protein of 185kDa, which is identified as c-erbB-2/HER-2/neu. Its

epitope is localized in the extracellular domain. C-erbB-2/HER-2 is a member of the EGFR family. This MAb is specific and shows minimal cross-reaction with other members of the EGFR-family. Receptors of this family are located on the plasma membrane and consist of an extracellular ligand-binding domain that is connected to a large intracellular domain by a single transmembrane sequence. c-erbB-2/HER-2 protein is over-expressed in a variety of carcinomas especially

those of breast and ovary.

Species/Host Mouse

Reactivity Human

**Conjugation** Unconjugated

**Tested Applications** IHC-P

Immunogen A recombinant human HER2 protein fragment within amino acids 311-462 was

used as the immunogen for the HER2 antibody. Its epitope is localized in the

extracellular domain.

**Preservatives** 0.2 mg/ml with 0.1 mg/ml rAlbumin (US sourced), 0.05% sodium azide

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

**Note** For research use only

**Application notes** Optimal dilution of the antibody should be determined by the researcher.

**Formula** 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide

**Isotype** Mouse IgG2b, kappa

**Clonality** Monoclonal

Clone Number ERBB2/3078






**Purity** Protein G affinity chromatography

Uniprot ID P04626


**Hazard Information** This HER2 antibody is available for research use only.

**Dilution Range** Immunohistochemistry (FFPE): 1-2ug/ml

**Expiration Date** 12 months from date of receipt.



IHC staining of FFPE human breast carcinoma with HER2 antibody (clone ERBB2/3078). HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.




IHC staining of FFPE human breast carcinoma with HER2 antibody (clone ERBB2/3078). HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>









IHC staining of FFPE human breast carcinoma with HER2 antibody (clone ERBB2/3078). HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.

## Human Protein Microarray Specificity Validation



Analysis of HuProt (TM) microarray containing more than 19000 full-length human proteins using HER2 antibody (clone ERBB2/3078). These results demonstrate the foremost specificity of the ERBB2/3078 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt (TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt (TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free HER2 antibody (clone ERBB2/3078) as confirmation of integrity and purity.

## **Biorbyt Ltd.**