

Product Datasheet

CD30 Antibody Cocktail (orb317428)

Description

Recognizes a single chain glycoprotein of 105/120kDa,


Species/Host

Mouse

Reactivity


Human

Conjugation

Other

Tested Applications

IHC-P



IHC:
Formalin-fixed,
paraffin-embedded
h...

Immunogen

A Co cell line established from a patient with Hodgkin's disease of T-cell lineage (Ber-H2) and human recombinant protein (CD30/412) were used as the immunogen for the CD30 antibody cocktail.

Preservatives

Bioreactor concentrate with 0.05% sodium azide

Storage

Store the CD30 antibody cocktail at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

Note

For research use only

Application notes

Optimal dilution of the CD30 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Formula

Bioreactor concentrate with 0.05% sodium azide

Isotype

Mouse IgGs

Clonality

Monoclonal

Purity

Unpurified high titer supernatant

Uniprot ID

P28908

Hazard Information

This CD30 antibody cocktail is available for research use only.

Dilution Range

Immunohistochemistry (FFPE): 1:100-1:200 for 30 min at RT (1)

Expiration Date

12 months from date of receipt.