

---

## Product Datasheet


### **N-Acylmannosamine-1-Dehydrogenase antibody (Biotin) (orb344308)**

**Description**


N-Acylmannosamine-1-Dehydrogenase



<b>Species/Host</b>	Goat
<b>Reactivity</b>	Bacteria
<b>Conjugation</b>	Biotin
<b>Tested Applications</b>	ELISA, IP, WB
<b>Immunogen</b>	N-Acyl Mannosamine-1-Dehydrogenase [Recombinant Sequence from Pseudomonas expressed in E.coli ]
<b>Preservatives</b>	0.01% (w/v) Sodium Azide
<b>Form/Appearance</b>	Lyophilized
<b>Concentration</b>	1.0 mg/mL
<b>Storage</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Note</b>	For research use only
<b>Application notes</b>	Anti-N-Acylmannosamine-1-Dehydrogenase antibody has been tested by western blot and is suitable to be assayed against 1.0 µg of N-Acylmannoseamine-1-Dehydrogenase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:7,000 to 1:35,000 of the reconstitution concentration is suggested for this product.
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Anti-N-Acylmannosamine-1-Dehydrogenase antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis



Biorbyt Goat anti N-Acylmannosamine-1-Deh...



Western Blot of Goat anti-N-Acylmannosea...