

Product Datasheet

Arginase antibody (Peroxidase) (orb344868)



www.biorbyt.com

Descriptionnts. Arginase antibody (Peroxidase)

Species/Host Rabbit

Reactivity Bovine

Conjugation HRP

Tested DOT, ELISA, WB

Applications

Immunogen Arginase [Bovine Liver]

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium

Azide!

Form/Appearance Liquid (sterile filtered)

Concentration 1 mg/ml

Storage Store vial at -20° C or below prior to opening. This vial

contains a relatively low volume of reagent (25 μ L). To minimize loss of volume dilute 1:10 by adding 225 μ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate

dilution when calculating final dilutions as

recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

Note For research use only

Application notes Anti-Arginase Peroxidase Antibody has been tested by

dot blot and is suitable to be assayed against 1.0 ug of Arginase [Bovine Liver] in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration

is suggested for this product.

Isotype IgG

Clonality Polyclonal

Purity Arginase is an IgG fraction antibody purified from

monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum as well as purified and partially purified Arginase [Bovine Liver]. Cross reactivity against Arginase from other tissues and species may occur but have not been

specifically determined



Dot Blot results of Rabbit Anti-Arginase...