


---

## Product Datasheet

### Beta galactosidase antibody (Biotin) (orb344931)

<b>Description</b>	Beta galactosidase antibody (Biotin)	
<b>Species/Host</b>	Rabbit	
<b>Conjugation</b>	Biotin	
<b>Tested Applications</b>	ELISA, IHC, WB	
<b>Immunogen</b>	Beta Galactosidase (E.coli)	Western Blot of Rabbit anti-Beta Galacto...
<b>Preservatives</b>	0.01% (w/v) Sodium Azide	
<b>Form/Appearance</b>	Liquid (sterile filtered)	
<b>Concentration</b>	1.0 mg/mL	
<b>Storage</b>	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.	
<b>Note</b>	For research use only	
<b>Application notes</b>	Anti-Beta Galactosidase Biotin Conjugated Antibody has been tested by Western blot and is suitable for ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. The antibody recognizes both frozen tissue sections, paraffin embedded tissue and 4% paraformaldehyde fixed tissue for most immunohistochemical analysis. A 1:5,000 dilution has been reported to be successful for staining by immunoblot of beta-galactosidase fusion proteins after transfer using a semi-dry transfer apparatus. A 1:1,500 dilution has been reported to detect beta-galactosidase in adult rat spinal cord tissue after infection with helper-dependent adenovirus expressing lacZ. In this particular experiment, tissue was perfused with 4% paraformaldehyde and cryostat-cut (35 µm) to produce free-floating sections. A 1:5,000 dilution has been reported to be successful for staining of brain sections from transgenic mice expressing nuclear beta-galactosidase when assayed by immunofluorescence microscopy. A 1:5,000 dilution has been reported for immunofluorescent staining of methanol fixed, devitellinized Drosophila embryos. Although a wide	

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive<br>Research Triangle Park<br>Durham, North Carolina<br>27709, United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558