

Product Datasheet

GFP antibody (Peroxidase) (orb345312)



www.biorbyt.com

Descriptionnts. GFP antibody (Peroxidase)

Species/Host Goat

Reactivity Other

Conjugation HRP

Tested ELISA, IHC, WB

Applications

Immunogen Recombinant Green Fluorescent Protein (GFP) fusion

protein corresponding to the full length amino acid sequence (246aa) derived from the jellyfish Aequorea

victoria.

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium

Azide!

Form/Appearance Lyophilized

Concentration 1.0 mg/mL

Storage Store GFP antibody at 4° C prior to restoration. For

extended storage aliquot contents and freeze at -20° C

or below. Avoid cycles of freezing and thawing.

Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Note For research use only

Application notes Polyclonal anti-GFP is designed to detect GFP and its

variants. Anti-GFP Peroxidase conjugated antibody has been tested by ELISA to detect GFP by ELISA (sandwich

or capture) for the direct binding of antigen and

recognizes wild type, recombinant and enhanced forms of GFP and by western blot. Biotin conjugated polyclonal anti-GFP used in a sandwich ELISA is well suited to

titrate GFP in solution using this antibody in

combination with Rockland's monoclonal anti-GFP (600-

301-215) using either form of the antibody as the capture or detection antibody. However, use the monoclonal form only for the detection of wild type or recombinant GFP as this form does not sufficiently detect 'enhanced' GFP. The detection antibody is typically conjugated to biotin and subsequently reacted with streptavidin conjugated HRP (code # S000-03). Fluorochrome conjugated polyclonal anti-GFP can be used to detect GFP by immunofluorescence microscopy

used to detect GFP by immunofluorescence microscopy in prokaryotic (E.coli) and eukaryotic (CHO cells) expression systems and can detect GFP containing inserts. Significant amplification of signal is achieved using fluorochrome conjugated polyclonal anti-GFP

ina 1 2 3 4 5 6 7 8 9 10 11 12 13 14 ina

Western Blot of Goat anti-GFP antibody P...