

Product Datasheet

HDAC-1 antibody (orb345508)



Description hts. HDAC-1 antibody

Species/Host Rabbit

Reactivity Human

Conjugation Unconjugated

Tested ELISA, IF, IHC, Multiplex Assay, WB

Applications

Immunogen Anti-HDAC-1 antibody was prepared from whole

rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a C-Terminal region near amino acids 450-482 of Human

HDAC-1.

Preservatives 0.01% (w/v) Sodium Azide

Form/Appearance Liquid (sterile filtered)

Concentration 1.33 mg/mL

Storage Store vial at -20° C prior to opening. Aliquot contents

and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior

to immediate use.

Note For research use only

Application notes Anti-HDAC-1 Antibody has been tested for use in

ELISA, immunohistochemistry, immunofluorescence, and western blot. Specific conditions for reactivity should be optimized by the end user. Specific nuclear staining is observed by IHC. Expect bands at 65 kDa in size corresponding to HDAC-1 by western blotting

in the appropriate cell lysate or extract.

Isotype IgG

Clonality Polyclonal

Purity Anti-HDAC-1 antibody is directed against human


HDAC-1 protein. HDAC-1 antibody was affinity

purified from monospecific antiserum by

immunoaffinity purification. A BLAST analysis was used to suggest reactivity with this protein from human, mouse, rat and chimpanzee sources based on 100% homology for the immunogen sequence. Cross reactivity may occur with HDAC-1 from bovine (82% homology) and chicken (80% homology) sources. Cross reactivity with HDAC-1 homologues

from other sources has not been determined

www.biorbyt.com



Immunofluorescence Microscopy of Rabbit

...





Immunofluorescence Microscopy of Rabbit

...



Immunohistochemistry of Rabbit Anti-HDAC...