
Product Datasheet

F(ab')₂ Rabbit IgG F(ab')₂ antibody (orb346790)

Excipients: Biotinylated.

F(ab')₂ Rabbit IgG F(ab')₂ antibody

Description	F(ab') ₂ Rabbit IgG F(ab') ₂ antibody
Species/Host	Goat
Reactivity	Rabbit
Conjugation	Unconjugated
Tested Applications	ELISA, IHC, WB
Immunogen	Rabbit IgG F(ab') ₂ fragment
Preservatives	None. 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Form/Appearance	Lyophilized
Concentration	10 mg/mL
Storage	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note	For research use only
Application notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 ⁶ cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.
Isotype	IgG F(ab') ₂
Clonality	Polyclonal
Purity	This product is a F(ab') ₂ fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt