
Product Datasheet

F(ab')₂ Rat IgG (H&L) antibody (orb346794)

Description

 F(ab')₂ Rat IgG (H&L) antibody

Species/Host

Rabbit

Reactivity

Rat

Conjugation

Unconjugated

Tested Applications

ELISA, IHC, WB

Immunogen

Rat IgG whole molecule

Preservatives

None

Form/Appearance

Lyophilized

Concentration

10.0 mg/mL

Storage

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Note

For research use only

Application notes

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10⁶ cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.

Isotype

 IgG F(ab')₂
Clonality

Polyclonal

Purity

This product is a F(ab')₂ fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion.