

Product Datasheet

Chicken IgG F(c) antibody (FITC) (orb346910)



www.biorbyt.com

kDa M 1 2 3 4 5

Western Blot of

Unconjugated Anti-Chicke...

Descriptionnts. Chicken IgG F(c) antibody (FITC)

Species/Host Rabbit

Reactivity Gallus

FITC Conjugation

Tested FC. FLISA. IF

Applications

Chicken IgG F(c) fragment **Immunogen**

0.01% (w/v) Sodium Azide **Preservatives**

Form/Appearance Lyophilized

Concentration 1.5 mg/mL

Store vial at 4° C prior to restoration. For extended Storage

> storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Note For research use only

Application notes Suitable for immunomicroscopy and flow cytometry or

FACS analysis as well as other antibody based

fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot

consistency, high titer and specificity.

Isotype IgG

Clonality Polyclonal

Purity This product was prepared from monospecific

> antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Chicken IgG, Chicken IgG F(c) and Chicken Serum. No reaction was observed against

Chicken IgG F(ab).

Dilution Range FLISA: 1:10,000 - 1:50,000, FC: 1:500 - 1:2,500, IF:

1:1,000 - 1:5,000

Expiration Date 12 months from date of receipt.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom