

Product Datasheet

Goat IgG F(ab')2 Antibody Peroxidase Conjugated Pre-Adsorbed (orb346995)

Description Goat IgG F(ab')2 antibody (Peroxidase)

Species/Host Rabbit

Reactivity Goat

Conjugation HRP

Tested Applications ELISA, IHC, WB

Immunogen Goat IgG F(ab')2 fragment

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Form/Appearance Lyophilized

Concentration 1.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Note For research use only

Application notes Anti-Goat IgG F(ab')2 Peroxidase Antibody has been tested by ELISA. This

product has been assayed against 1.0 μ g of Goat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:8,000 to 1:62,000 of the reconstitution concentration is suggested for this

product.

Isotype IgG





Clonality Polyclonal

Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Goat IgG coupled to agarose beads followed by solid


phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-

Peroxidase, anti-Rabbit Serum, Goat IgG, Goat IgG F(ab')2 and Goat Serum. No

reaction was observed against Goat IgG F(c) or Human Serum Proteins.

Dilution Range ELISA: 1:10,000 - 1:50,000, IHC: 1:500 - 1:2,500, WB: 1:1,000 - 1:10,000

Expiration Date 12 months from date of receipt.



ELISA results of purified Rabbit Anti-Goat IgG F(ab')2 Antibody Peroxidase Conjugated tested against purified Goat IgG F(ab')2. Each well was coated in duplicate with 1.0 μ g of Goat IgG F(ab')2. The starting dilution of antibody was 5 μ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558