

Product Datasheet

Human IgG (H&L) Antibody Peroxidase Conjugated (orb347222)

Description Human IgG (H&L) antibody (Peroxidase)

Species/Host Goat

Reactivity Human

Conjugation HRP

Tested Applications DOT, ELISA, IHC, WB

Immunogen Anti-Human IgG (H&L) was produced by repeated immunization with human IgG

whole molecule in goat.

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Form/Appearance Lyophilized

Concentration 2.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Note For research use only

Application notes Anti-Human IgG (H&L) peroxidase conjugated antibody has been tested by dot

blot and ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as

other peroxidase-antibody based enzymatic assays requiring lot-to-lot

consistency.

Isotype IgG

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>





Clonality

Polyclonal

Purity


This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG and Human Serum.

Dilution Range


ELISA: 1:20,000 - 1:80,000, IHC: 1:1,000 - 1:5,000, WB: 1:5,000 - 1:25,000

Expiration Date

12 months from date of receipt.



Dot Blot of Human IgG. Lane 1: 100 ng. Lane 2: 33.3 ng. Lane 3: 11.1 ng. Lane 4: 3.7 ng. Lane 5: 1.23 ng. Secondary Antibody: orb347222 HRP Gt-a-Hu IgG 1:1000. Blocking Buffer: BlockOut orb348644 for 1 hour RT.



ELISA results of purified Goat anti-Human IgG Antibody Peroxidase conjugated tested against BSA-conjugated peptide of immunizing peptide. Each well was coated in duplicate with 1.0 μ g of Human IgG (p/n orb346219). The starting dilution of antibody was 5 μ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% fish gelatin as blocking buffer and TMB substrate p/n orb348651.