

## **Product Datasheet**

## Human IgG F(ab')2 Antibody Biotin Conjugated (orb347243)

**Description** Human IgG F(ab')2 antibody (Biotin)

Species/Host Goat

**Reactivity** Human

**Conjugation** Biotin

**Tested Applications** ELISA, IHC, WB

**Immunogen** Human IgG F(ab')2 fragment

**Preservatives** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Form/Appearance Lyophilized

**Concentration** 2.0 mg/mL

**Storage** Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

**Note** For research use only

**Application notes** Anti-Human IgG F(ab')2 Biotin Antibody has been tested by ELISA and is ideal for

ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow

Cytometry as well as other antibody detection methods.

**Isotype** IgG

**Clonality** Polyclonal





**Purity** 


This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Human IgG, Human IgG F(ab')2 and Human Serum. No reaction was observed against Human IgG F(c).

**Dilution Range** 

ELISA: 1:20,000 - 1:500,000, IHC: 1:1,000 - 1:5,000, WB: 1:2,000 - 1:10,000

**Expiration Date** 

12 months from date of receipt.



ELISA of purified Goat Anti-Human IgG F(ab')2 Antibody Biotin Conjugated tested against purified Human IgG F(ab')2 Biotin. Each well was coated in duplicate with 1.0  $\mu$ g of Human IgG F(ab')2 (p/n orb346221). The working dilution is 1:108000. The starting dilution of antibody was 5  $\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using HRP Conjugate Stabilizer, Streptavidin-HRP conjugated (p/n orb348767) and TMB substrate (p/n orb348651).