

Product Datasheet

Human IgG (H&L) Antibody Peroxidase Conjugated (orb347290)

Description Human IgG (H&L) antibody (Peroxidase)

Species/Host Rabbit

Reactivity Human

Conjugation HRP

Tested Applications DOT, ELISA, IHC, WB

Immunogen Anti-Human IgG whole molecule was produced by repeated immunization with

Human IgG whole molecule in rabbit.

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Form/Appearance Lyophilized

Concentration 2.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Note For research use only

Application notes Anti-Human IgG (H&L) Peroxidase Conjugated Antibody has been tested by ELISA

and Dot Blot and is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as

other peroxidase-antibody based enzymatic assays requiring lot-to-lot

consistency.

Isotype IgG

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>





Clonality Polyclonal


Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Human IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-


Peroxidase, anti-Rabbit Serum, Human IgG and Human Serum.

Dilution Range ELISA: 1:150,000 - 1:250,000, IHC: 1:1,000 - 1:5,000, WB: 1:5,000 - 1:25,000

Expiration Date 12 months from date of receipt.



Dot Blot Results of Rabbit Anti-Human IgG HRP Conjugated Antibody. Human IgG sample loaded 1- 100 ng, 2- 33.3 ng, 3- 11.1 ng, 4- 3.7 ng, 5- 1.2 ng. Rabbit Anti-Human IgG Peroxidase Conjugated Antibody 1:1000 for 60 mins at RT. Blocking Buffer for Fluorescent Western Blotting (p/n orb348637) for 1hr at RT.



ELISA results of Rabbit anti-Human IgG Antibody Peroxidase Conjugated tested against purified Human IgG protein. Each well was coated in duplicate with 1.0 μ g of Human IgG (p/n orb346219). The starting dilution of antibody was 5 μ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% fish gelatin as blocking buffer and TMB substrate p/n orb348651.