

---

## Product Datasheet

### Mouse IgG (H&L) antibody (RPE) (orb347406)

**Description**

Mouse IgG (H&amp;L) antibody (RPE)

**Species/Host**

Goat

**Reactivity**


Mouse

**Conjugation**

RPE

**Tested**

DOT, FC, IF, WB

**Applications**

 Dot Blot of  
Goat anti-  
Mouse IgG  
Antibody...

**Immunogen**

Anti-Mouse IgG was produced by repeated immunization with Mouse IgG whole molecule in goat.

**Preservatives**

0.01% (w/v) Sodium Azide

**Form/Appearance**

Lyophilized

**Concentration**

0.5 mg/mL

**Storage**

Store vial at 4° C prior to restoration. Restore with deionized water (or equivalent). This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Centrifuge product if not completely clear after standing at room temperature. Do not freeze after reconstitution. Store reagent in the dark. Use subdued lighting during handling and incubation of cells prior to analysis.

**Note**

For research use only

**Application notes**

Anti-Mouse IgG (H&L) Phycoerythrin conjugated antibody has been tested by western blot and dot blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.

**Isotype**

IgG

**Clonality**

Polyclonal

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, conjugation to phycoerythrin and subsequent purification, and extensive dialysis against