

## Product Datasheet

# Mouse IgM (mu chain) Antibody Peroxidase Conjugated (orb347431)

<b>Description</b>	Peroxidase Conjugated Mouse IgM (Mu Chain) Antibody
<b>Species/Host</b>	Goat
<b>Reactivity</b>	Mouse
<b>Conjugation</b>	HRP
<b>Tested Applications</b>	ELISA, IHC, WB
<b>Immunogen</b>	Mouse IgM whole molecule
<b>Preservatives</b>	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
<b>Form/Appearance</b>	Lyophilized
<b>Concentration</b>	1.0 mg/mL
<b>Storage</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Note</b>	For research use only
<b>Application notes</b>	Mouse IgM (mu chain) peroxidase conjugated Antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, lot-to-lot consistency, high titer and specificity.
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal

**Biorbyt Ltd.**

7 Signet Court, Swann's Road,  
Cambridge, CB5 8LA, United Kingdom  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)


**Biorbyt LLC.**

68 TW Alexander Drive,  
Durham, NC, 27713, United States  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)


**Purity** This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgM and Mouse Serum. No reaction was observed against other mouse heavy or light chain proteins.

**Dilution Range** ELISA: 1:30,000, IHC: 1:500 - 1:2,000, WB: 1:1,000 - 1:5,000

**Expiration Date** 12 months from date of receipt.



ELISA results using Goat Anti-Mouse IgM HRP. Determination of mAb titer and affinity. (A) Titration of mAb by an indirect ELISA. The mAb was serially diluted in 1:3. The optimum working concentration was determined for a midpoint of the steep portion of the curve. (B) The measurement of antibody relative affinity by thiocyanate elution assay. The affinity index was estimated by the molarity of NH<sub>4</sub>SCN causing 50% reduction from initial absorbance in the elution curves. All experiments were carried out in triplicate and the results were calculated from three independent experiments.



Immunochemical Staining using Goat Anti-Mouse IgM HRP. Identification of intact *B. melitensis* strain by ICS with mAbs. The intact bacteria of *B. melitensis* strain were stained by ICS with individual mAbs. Saturated with goat anti-mouse IgG and IgM HRP conjugate. Bacteria were visualized with diaminobenzidine (DAB) substrate for color development. G-, Gram staining for bacterial control of *B. melitensis* strain examined under white light with a microbiological microscope.