




---

## Product Datasheet

### Rabbit IgG F(c) antibody (orb347664)

<b>Description</b>	Rabbit IgG F(c) antibody	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Rabbit	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	DOT, ELISA, IHC, WB	Dot Blot of Goat Anti-RABBIT IgG F(c) An...
<b>Immunogen</b>	Anti Rabbit IgG F(c) was produced by repeated immunization with rabbit IgG F(c) fragment in goat.	
<b>Preservatives</b>	0.01% (w/v) Sodium Azide	
<b>Form/Appearance</b>	Liquid (sterile filtered)	
<b>Concentration</b>	2.7mg/mL	
<b>Storage</b>	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.	
<b>Note</b>	For research use only	Dot Blot Results of Goat Anti-Rabbit F(c)...
<b>Application notes</b>	Anti-Rabbit IgG F(c) is tested by immunoelectrophoresis, ELISA, dot blot, and western blot. This product is suitable for use in western-blot, competitive western-blot, and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user. Expect ~25kDa for Rabbit IgG F(c) Fragment.	
<b>Isotype</b>	IgG	
<b>Clonality</b>	Polyclonal	ELISA results of purified Goat Anti-Rabb...
<b>Purity</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab) or Human Serum Proteins.	
<b>Dilution Range</b>	ELISA: 1:20,000 - 1:100,000, IHC: 1:1,000 - 1:5,000, WB: 1:2,000 - 1:10,000	
<b>Expiration Date</b>	12 months from date of receipt.	