

Product Datasheet

Rabbit IgG F(c) antibody (Peroxidase) (orb347674)



Descriptionnts.

www.biorbyt.com

KDa M 1 2 3 4 5

Species/Host Goat

Reactivity Rabbit

Conjugation HRP

Tested ELISA, IHC, WB

Applications

Immunogen Anti-Rabbit IgG F(c) fragment was produced by

repeated immunization with Rabbit IgG F(c) fragment

in goat.

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium

Rabbit IgG F(c) antibody (Peroxidase)

Azide!

Form/Appearance Lyophilized

Concentration 2.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended

storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute

only prior to immediate use.

Note For research use only

Application notes Anti-Rabbit IgG Fc peroxidase conjugated antibody

has been tested by ELISA and western blot and is suitable for use in immunoelectrophoresis, westernblot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.

Isotype IgG

Clonality Polyclonal

Purity Anti-Rabbit IgG F(c) antibody was prepared from

monospecific antiserum by immunoaffinity

chromatography using Rabbit IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab).

Dilution Range ELISA: 1:110,000, IHC: 1:1,000 - 1:5,000, WB: 1:2,000

- 1:20,000

Expiration Date 12 months from date of receipt.

Western Blot of Anti-Rabbit IgG F(c) (GO...

iOa 1
150 - 1
100 - 80 - 60 - 50 - 40 - 30 - 20 - 20 -

Western blot of Peroxidase conjugated Go...