

Product Datasheet

Rat IgG (H&L) antibody (Peroxidase) (orb347892)



Descriptionnts.

www.biorbyt.com

Rat IgG (H&L) antibody (Peroxidase)

Species/Host Donkey

Reactivity Rat

Conjugation HRP

Tested ELISA, IHC, WB

Applications

Immunogen Rat IgG whole molecule

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium

Azide!

Form/Appearance Lyophilized

Concentration 1.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended

storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid.

Dilute only prior to immediate use.

Note For research use only

Application notes Anti-Rat IgG Peroxidase has been tested by ELISA and

is assayed against 1.0 ug of Rat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-

ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:15,000 to 1:60,000 is suggested

for this product.

Isotype IgG

Clonality Polyclonal

Purity This product was prepared from monospecific

antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted

reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Rat IgG and Rat Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and

Sheep Serum Proteins.

Dilution Range ELISA: 1:10,000 - 1:50,000, IHC: 1:500 - 1:2,500, WB:

1:1,000 - 1:10,000

ICu M 1 2 3 4 5

Western Blot of Anti-Rat IgG (H&L) (DONK...

Carolina < br > 27709. United States

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North