

Product Datasheet

F(ab')2 Mouse IgG F(c) antibody (FITC) (orb348247)



www.biorbyt.com

Descriptionnts. F(ab')2 Mouse IgG F(c) antibody (FITC)

Species/Host Goat

Reactivity Mouse

Conjugation FITC

Tested DOT, FC, FLISA, IF

Applications

Immunogen Mouse IgG F(c) fragment

Preservatives 0.01% (w/v) Sodium Azide

Form/Appearance Lyophilized

Concentration 1.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended

storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid.

Dilute only prior to immediate use.

Note For research use only

Application notes F(ab')2 Anti-Mouse IgG F(c) Fluorescein Antibody has

been tested by dot blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels,

absence of F(c) mediated binding, lot-to-lot

consistency, high titer and specificity. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial

platforms.

Isotype IgG F(ab')2

Clonality Polyclonal

Purity This product was prepared from monospecific

antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG, Mouse IgG

F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(ab) or Bovine, Horse and Human Serum Proteins.



Dot Blot results of Goat F(ab')2 Anti-Mo...

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North