

Product Datasheet

F(ab')₂ Mouse IgG (H&L) Antibody Peroxidase Conjugated (orb348250)

Description	F(ab') ₂ Mouse IgG (H&L) antibody (Peroxidase)
Species/Host	Goat
Reactivity	Mouse
Conjugation	HRP
Tested Applications	ELISA, IHC, WB
Immunogen	Mouse IgG whole molecule
Preservatives	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Form/Appearance	Lyophilized
Concentration	1.0 mg/mL
Storage	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note	For research use only
Application notes	F(ab') ₂ Anti-Mouse IgG (H&L) Peroxidase Antibody has been tested by ELISA and western blot and is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.
Isotype	IgG F(ab') ₂


Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Clonality	Polyclonal
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).
Dilution Range	ELISA: 1:10,000 - 1:50,000, IHC: 1:500 - 1:2,500, WB: 1:1,000 - 1:10,000
Expiration Date	12 months from date of receipt.



Western blot of Peroxidase conjugated Goat F(ab')₂ Anti-Mouse IgG secondary antibody. Lane 1: Mouse IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Peroxidase goat secondary antibody at 1:1000 for 60 min at RT. Blocking: orb348637 for 30 min at RT. Observed/Predicted size: 25 & 55 kDa, 25 & 55 kDa for Mouse IgG. Other band(s): None.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)