

Product Datasheet

F(ab')2 Rabbit IgG F(ab')2 antibody (Peroxidase) (orb348309)



www.biorbyt.com

kDa 1

80 **–** 60 **–**

80-

Western Blot of Goat anti-F(ab')2 Rabbit...

Descriptionnts. F(ab')2 Rabbit IgG F(ab')2 antibody (Peroxidase)

Species/Host Goat

Reactivity Rabbit

Conjugation HRP

Tested ELISA, IHC, WB

Applications

Immunogen Anti-Rabbit IgG F(ab')2 Antibody was produced in

goat by repeated immunization with Rabbit IgG

F(ab')2 fragment.

Preservatives 0.01% (w/v) Gentamicin Sulfate. Do NOT add

Sodium Azide!

Form/Appearance Lyophilized

Concentration 1.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended

storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Note For research use only

Application notes F(ab')2 Anti-Rabbit IgG F(ab')2 Peroxidase

Antibody has been tested by ELISA and western blot and is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels,

absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

Isotype IgG F(ab')2

Clonality Polyclonal

Purity Anti-Rabbit IgG F(ab')2 Antibody was prepared

from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to

agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serum. No reaction was observed against

Carolina < br > 27709. United States