

Product Datasheet

F(ab')2 Rabbit IgG (H&L) antibody (TRITC) (orb348320)



www.biorbyt.com

Descriptionnts. F(ab')2 Rabbit IgG (H&L) antibody (TRITC)

Species/Host Donkey

Reactivity Rabbit

Conjugation TRITC

Tested IF, WB

Applications

Immunogen Rabbit IgG whole molecule

Preservatives 0.02 M Potassium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Form/Appearance Lyophilized

Concentration 1.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended

storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Note For research use only

Application notes F(ab')2 Anti-Rabbit IgG Rhodamine Conjugated

Antibody has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high

titer and specificity.

Isotype IgG F(ab')2

Clonality Polyclonal

Purity This product was prepared from monospecific

antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Bovine, Chicken, Goat, Guinea

Western Blot of Donkey anti-Rabbit IgG P...