
Product Datasheet

**F(ab')₂ Rat IgG F(c) antibody (Peroxidase)
(orb348343)**

Description

 F(ab')₂ Rat IgG F(c) antibody (Peroxidase)

Anti-Rat IgG F(c) Sensitivity

Species/Host

Goat

Reactivity


Rat

Conjugation

HRP

Tested

ELISA, IHC, WB

Applications


ELISA results of purified F(ab')₂ Goat a...

Immunogen

Rat IgG F(c) fragment

Preservatives

0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Form/Appearance

Lyophilized

Concentration

1.0 mg/mL

Storage

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Note

For research use only

Application notes

F(ab')₂ Anti-Rat IgG F(c) Peroxidase Antibody has been tested by ELISA and is assayed against 1.0 ug of Rat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:200,000 of the reconstitution concentration is suggested for this product for ELISA and Western blots and a 1:500 to 1:5,000 for immunohistochemistry on tissue sections.

Isotype

 IgG F(ab')₂
Clonality

Polyclonal

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rat IgG, Rat IgG F(c) and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat IgG F(ab) or Bovine, Horse, Human Serum Proteins.