

Product Datasheet

Fab Goat IgG (H&L) antibody (FITC) (orb348429)



www.biorbyt.com

Descriptionnts. Fab Goat IgG (H&L) antibody (FITC)

Species/Host Donkey

Reactivity Goat

FITC Conjugation

Tested FC. FLISA. IF. WB

Applications

Goat IgG whole molecule **Immunogen**

0.01% (w/v) Sodium Azide **Preservatives**

Form/Appearance Lyophilized

Concentration 1.0 mg/mL

Store vial at 4° C prior to restoration. For extended Storage

> storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Note For research use only

Application notes Fab Anti-Goat IgG Fluorescein Antibody has been

tested by western blot and is suitable for

immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels,

absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

Isotype IgG Fab

Clonality Polyclonal

Purity This product was prepared from monospecific

> antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Donkey Serum. No reaction was observed against

anti-papain or anti-Donkey IgG F(c).

Dilution Range FLISA: 1:10,000 - 1:50,000, FC: 1:500 - 1:2,500, IF:

1:1,000 - 1:5,000

Expiration Date 12 months from date of receipt.

Western blot of

Fluorescein conjugated D...