

Product Datasheet

Anti-IL-2R alpha (CD25) [Basiliximab] (orb348865)

Description Human monoclonal antibody to CD25

Species/Host Human

Reactivity Human, Primate

Conjugation Unconjugated

Tested Applications Blocking, FC, IF, IHC, WB

Immunogen Basiliximab was prepared by immunizing BALB/c mice with CTC 16 cells.

Target IL-2R alpha (CD25)

Preservatives PBS with 0.02% Proclin 300.

Concentration 1 mg/ml

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only

Isotype Human IgG1

Clonality Monoclonal

Clone Number Basiliximab


Purity Purified

Uniprot ID P01589


Expiration Date 12 months from date of receipt.








Flow-cytometry using anti-CD25 antibody Basiliximab Cynomolgus monkey lymphocytes were stained with an isotype control (orb256390, panel A) or the rabbit-chimeric version of Basiliximab (orb348908, panel B) at a concentration of 1 μ g/ml for 30 mins at RT. After washing, bound antibody was detected using a AF488 conjugated donkey anti-rabbit antibody and cells analysed on a FACSCanto flow-cytometer.




Flow-cytometry using anti-CD25 antibody Basiliximab Rhesus monkey lymphocytes were stained with an isotype control (orb256390, panel A) or the rabbit-chimeric version of Basiliximab (orb348908, panel B) at a concentration of 1 μ g/ml for 30 mins at RT. After washing, bound antibody was detected using a AF488 conjugated donkey anti-rabbit antibody and cells analysed on a FACSCanto flow-cytometer.




Flow-cytometry using the anti-CD25 (IL2R) research biosimilar antibody Basiliximab Human lymphocytes were stained with an isotype control (orb256390, panel A) or the rabbit-chimeric version of Basiliximab (orb348908, panel B) at a concentration of 1 μ g/ml for 30 mins at RT. After washing, bound antibody was detected using a AF488 conjugated donkey anti-rabbit antibody and cells analysed on a FACSCanto flow-cytometer.








Immunofluorescent staining of rat thymus using anti-CD25 antibody. Formaldehyde-fixed rat thymus slices were stained with orb348865 at 5 μ g/ml and detected with a FITC-conjugated secondary antibody.



Immunohistochemical staining of rat spleens using anti-CD25 antibody. Formalin fixed rat spleen slices were were stained with a orb348865 at 5 μ g/ml. The anti-human CD25 antibody also shows some cross-reactivity with rat in IHC.



Western Blot using anti-IL2R alpha (CD25) antibody. Human thyroid cancer (lane 1), rat spleen (lane 2), mouse lymph node (lane 3), mouse thymus (lane 4) and rat small intestines (lane 5)samples were resolved on a 10% SDS PAGE gel and blots probed with orb348865 at 1 μ g/ml before being detected by a secondary antibody.

 $\begin{aligned} & \text{Email: } \underline{\text{info@biorbyt.com}}, \ \underline{\text{support@biorbyt.com}} \\ & \text{Phone: } \underline{+1 \ (415) \ 906\text{-}5211} \ \big| \ \text{Fax: } \underline{+1 \ (415) \ 651\text{-}8558} \end{aligned}$