

# **Product Datasheet**

# Anti-TGF beta Receptor I/TGFBR1 Antibody (orb381103)

Description	Anti-TGF beta Receptor I/TGFBR1 Antibody. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Species/Host	Rabbit
Reactivity	Human, Mouse, Rat
Conjugation	Unconjugated
Tested Applications	WB
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human TGFBR1, identical to the related mouse and rat sequences.
Form/Appearance	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 $\mu$ g/ml.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at - 20°C in small aliquots to prevent freeze-thaw cycles.
Note	For research use only
Application notes	Western blot, $0.1$ - $0.5\mu$ g/ml, Human, Mouse, Rat. Add $0.2$ ml of distilled water will yield a concentration of 500ug/ml
lsotype	Rabbit IgG
Clonality	Polyclonal
Antibody Type	Primary Antibody
MW	56 kDa
Uniprot ID	P36897

### **Biorbyt Ltd.**

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0) 1223 859-353</u> | Fax: <u>+1 (415) 651-8558</u>


### **Biorbyt LLC.**

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>



## **Biorbyt.com**

## **Expiration Date**



12 months from date of receipt.

Western blot analysis of TGFBR1 using anti-TGFBR1 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat cardiac muscle tissue lysates, Lane 2: mouse liver tissue lysates, Lane 3: HELA whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TGFBR1 antigen affinity purified polyclonal antibody at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for TGFBR1 at approximately 56 kDa. The expected band size for TGFBR1 is at 56 kDa.

#### **Biorbyt Ltd.**

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558

### **Biorbyt LLC.**

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>