


Product Datasheet

Hdac2 antibody (orb420062)


| | |
|----------------------------|--|
| Description | Goat polyclonal antibody to Hdac2 |
| Species/Host | Goat |
| Reactivity | Bovine, Canine, Human, Mouse, Porcine, Rat |
| Conjugation | Unconjugated |
| Tested Applications | ELISA, FC, IF, WB |
| Target | Hdac2 (mouse) |
| Preservatives | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Storage | Aliquot and store at -20°C. Minimize freezing and thawing. |
| Note | For research use only |
| Application notes | WB: Approx 55-60kDa band observed in nuclear lysates of cell line HEK293 (calculated MW of 55.3kDa according to NP_032255.2). Recommended concentration: 1-2 µg/ml. Primary incubation 1 hour at room temperature. |
| Protein Sequence | PEDAVHEDSGDE |
| Clonality | Polyclonal |
| MW | 55.3 |
| Entrez | 3066 |
| Dilution Range | ELISA: 1:2000, WB: 1-2 µg/ml, IF/ICC: 10µg/ml, FACS: 10ug/ml |
| Expiration Date | 12 months from date of receipt. |



Western blot analysis of HEK293 nuclear ...



Flow cytometric analysis of HeLa cells u...



Immunofluorescence analysis of NIH3T3 ce...