

Product Datasheet

Mouse IgG1 Isotype Control antibody (orb43531)



www.biorbyt.com

Descriptionnts. Mouse IgG1 Isotype Control

Conjugation Unconjugated

Tested ELISA, FC, ICC, IHC-Fr, IHC-P, IP, WB

Applications

Preservatives Phosphate buffered saline (PBS), pH 7.4, 15 mM

sodium azide

Concentration 1 mg/ml

Storage Store at 2-8°C. Do not freeze.

Note For research use only

Application notes Negative control: The reagent is intended as an

isotype control to establish the amount of nonspecific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above

 $10~\mu g/ml$ of incubation mixture), change the conditions of your experiment to reduce the

background.

Isotype Mouse IgG1

Clonality Monoclonal


Purity Purified by protein-A affinity chromatography.

Dilution Range Negative control: The reagent is intended as an

isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above

 $10 \mu g/ml$ of incubation mixture), change the conditions of your experiment to reduce the

background.



Flow cytometry surface nonspecific stain...