
Product Datasheet

Cytokeratin 10/13 antibody (orb43753)

Description

Mouse Monoclonal to Cytokeratin 10/13.


Reactivity


Feline, Human

Conjugation

Unconjugated

Tested Applications

IHC-Fr, IHC-P, WB


Immunogen

Cytoskeletal preparation extracted from human ectocervical epithelium.

Immunohistochemistry staining of tonsil ...

Target

Cytokeratin 10/13

Preservatives

Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Concentration

1 mg/ml

Storage

Store at 2-8°C. Do not freeze.

Note

For research use only

Application notes

Immunohistochemistry (paraffin sections): Standard ABC technique (DAB+), recommended dilution: 10 µg/ml (1:100), positive tissue: tonsil. Pretreatment: 0.1% pepsin in 0.1 M HCl for 30 min at room temperature or Target Retrieval Solution - High pH. The antibody DE-K13 reacts exclusively with Cytokeratin 13. Immunohistochemistry (frozen sections): Use at an assay dependent dilution. Western blotting: recommended dilution: 1-3 µg/ml.

Isotype

Mouse IgG2a

Clonality

Monoclonal

Purity

Purified by protein-A affinity chromatography.

Dilution Range

Immunohistochemistry (paraffin sections): Standard ABC technique (DAB+), recommended dilution: 10 µg/ml (1:100), positive tissue: tonsil. Pretreatment: 0.1% pepsin in 0.1 M HCl for 30 min at room temperature or Target Retrieval Solution - High pH. The antibody DE-K13 reacts exclusively with Cytokeratin 13. Immunohistochemistry (frozen sections): Use at an assay dependent dilution. Western blotting: recommended dilution: 1-3 µg/ml.

Expiration Date

12 months from date of receipt.