
Product Datasheet

FTL Antibody / Ferritin light chain (orb606893)

Description

Mammalian ferritins consist of 24 subunits made up of 2

Species/Host

Mouse

Reactivity

Human

Conjugation

Unconjugated

Tested Applications

IHC-P, WB

Immunogen

Amino acids 38-165 from the human protein were used as the immunogen for this FTL antibody.

Preservatives

0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin (US sourced) and 0.05% sodium azide

Storage

Store the FTL antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

Note

For research use only

Application notes

Optimal dilution of the FTL antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Formula

0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide

Isotype

Mouse IgG2a, kappa

Clonality

Monoclonal

Purity

Protein G affinity chromatography

Uniprot ID

[P02792](#)

Hazard Information


This FTL antibody is available for research use only.

Dilution Range

Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 0.1-0.2ug/ml for 30 min at RT

Expiration Date

12 months from date of receipt.




IHC testing of FFPE human prostate with ...



Western blot testing of human HeLa cell ...

Human Protein Microarray Specificity Validation



Analysis of HuProt (TM) microarray conta...