



## Product Datasheet

### c-Myc Antibody (orb639824)

<b>Description</b>	It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa ...
<b>Species/Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Tested Applications</b>	ELISA, FACS
<b>Immunogen</b>	Amino acids 408-439 were used as the immunogen for the recombinant c-Myc antibody. The specific epitope has been localized to amino acids 410-419 (EQKLISEEDL).
<b>Preservatives</b>	0.2 mg/ml with 0.1 mg/ml rAlbumin (US sourced), 0.05% sodium azide
<b>Storage</b>	Store the recombinant c-Myc antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).
<b>Note</b>	For research use only
<b>Application notes</b>	Optimal dilution of the antibody should be determined by the researcher.
<b>Formula</b>	0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
<b>Isotype</b>	Rabbit IgG, kappa
<b>Clonality</b>	Recombinant
<b>Purity</b>	Protein A affinity chromatography
<b>Uniprot ID</b>	<b>P01106</b>
<b>Hazard Information</b>	This recombinant c-Myc antibody is available for research use only.
<b>Dilution Range</b>	ELISA (order BSA-free format for coating), Flow cytometry: 1-2ug/million cells in 0.1ml
<b>Expiration Date</b>	12 months from date of receipt.



Flow cytometry testing of permeabilized ...



SDS-PAGE analysis of purified, BSA-free ...