

## **Product Datasheet**

## Anti-CD31/PECAM1 Antibody Picoband (monoclonal, 2D4) (orb654278)

**Description** Anti-CD31/PECAM1 Antibody (monoclonal, 2D4). Tested in IHC, IHC-F, WB

applications. This antibody reacts with Human.

Species/Host Mouse

**Reactivity** Human

**Conjugation** Unconjugated

**Tested Applications** IHC, IHC-Fr, WB

Immunogen E.coli-derived human CD31 recombinant protein (Position: Q28-G382). Human

CD31 shares 65% and 68% amino acid (aa) sequences identity with mouse and

rat CD31, respectively.

Form/Appearance Lyophilized

**Concentration** Adding 0.2 ml of distilled water will yield a concentration of 500 μg/ml.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

**Note** For research use only

**Application notes** Western blot, 0.1-0.5μg/ml, Human Immunohistochemistry (Paraffin-embedded

Section), 0.5-1µg/ml, Human Immunohistochemistry (Frozen Section), 0.5-1µg/ml, Human. Add 0.2ml of distilled water will yield a concentration of

500ug/ml

**Isotype** Mouse IgG2b

**Clonality** Monoclonal

Clone Number 2D4

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>







**MW** 130 kDa

Uniprot ID P16284

**Expiration Date** 12 months from date of receipt.




Western blot analysis of CD31/PECAM1 using anti-CD31/PECAM1 antibody (orb654278). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human Jurkat whole cell lysates; Lane 2: human placenta tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-CD31/PECAM1 antigen affinity purified monoclonal antibody (Catalog # orb654278) at 0.5 μg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # orb90502) with Tanon 5200 system. A specific band was detected for CD31/PECAM1 at approximately 130KD. The expected band size for CD31/PECAM1 is at 130KD.




IHC analysis of CD31/PECAM1 using anti-CD31/PECAM1 antibody (orb654278). CD31/PECAM1 was detected in paraffinembedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1µg/ml mouse anti-CD31/PECAM1 Antibody (orb654278) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # orb90443) with DAB as the chromogen.








IHC analysis of CD31/PECAM1 using anti-CD31/PECAM1 antibody (orb654278). CD31/PECAM1 was detected in paraffinembedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu$ g/ml mouse anti-CD31/PECAM1 Antibody (orb654278) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # orb90443) with DAB as the chromogen.



IHC analysis of CD31/PECAM1 using anti-CD31/PECAM1 antibody (orb654278). CD31/PECAM1 was detected in paraffinembedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu$ g/ml mouse anti-CD31/PECAM1 Antibody (orb654278) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # orb90443) with DAB as the chromogen.



IHC analysis of CD31/PECAM1 using anti-CD31/PECAM1 antibody (orb654278). CD31/PECAM1 was detected in frozen section of human placenta tissue. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu$ g/ml mouse anti-CD31/PECAM1 Antibody (orb654278) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # orb90443) with DAB as the chromogen.