

---

## Product Datasheet

### Goat anti-Rabbit IgG (H&L) antibody (TRITC) (orb700102)

**Description**

Goat polyclonal antibody against Goat

	TRITC
<b>Species/Host</b>	Goat
<b>Conjugation</b>	TRITC
<b>Form/Appearance</b>	Lyophilized
<b>Storage</b>	Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming.
<b>Note</b>	For research use only
<b>Application notes</b>	This antibody reacts with the heavy chains on rabbit IgG and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin rabbit serum proteins based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Immunogen affinity purified IgG.
<b>Dilution Range</b>	The optimal working dilution should be determined by the investigator, (ICC), immunohistochemistry ( IHC) (Flow cyt)
<b>Expiration Date</b>	12 months from date of receipt.