
Product Datasheet

**Goat anti-Bovine IgG (H&L) antibody (FITC)
(orb700452)**

Description

Goat polyclonal antibody against Goat

FITC

Species/Host

Goat

Reactivity

Bovine

Conjugation

FITC

Tested Applications

FC, IL

Immunogen

Purified Bovine IgG, whole molecule

Form/Appearance

Liquid

Storage

Store at -20° C.

Note

For research use only

Application notes

Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Bovine IgG (H&L). Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Based on IEP, this antibody reacts with:

- heavy (γ) chains on bovine IgG
- light chains on all bovine immunoglobulins.

Based on IEP, no reactivity is observed to:

- non-immunoglobulin bovine serum immunoglobulins.

Clonality

Polyclonal

Purity

Immunogen affinity purified goat IgG.

Uniprot ID

DAA31645

Dilution Range

1 : 20-1 : 2000 (Flow cys), (IL), The optimal working dilution should be determined by the investigator

Expiration Date

12 months from date of receipt.