

---

## Product Datasheet

**Goat anti-Donkey IgG (H&L) antibody (Biotin)  
(orb700465)**

**Description**

Goat polyclonal antibody against Goat

|                            |   |
|----------------------------|---|
|                            | Biotin  |
| <b>Species/Host</b>        | Goat  |
| <b>Reactivity</b>          | Donkey  |
| <b>Conjugation</b>         | Biotin  |
| <b>Tested Applications</b> | FC, IL  |
| <b>Immunogen</b>           | Purified Donkey IgG, whole molecule,  |
| <b>Form/Appearance</b>     | Liquid  |
| <b>Storage</b>             | Store at -20° C.  |
| <b>Note</b>                | For research use only   |
| <b>Application notes</b>   | Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Donkey IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: heavy (γ) chains on donkey IgG light chains on all donkey immunoglobulins. Based on IEP, no reactivity is observed to: non-immunoglobulin donkey serum immunoglobulins. |
| <b>Clonality</b>           | Polyclonal  |
| <b>Purity</b>              | Immunogen affinity purified goat IgG.   |
| <b>Dilution Range</b>      | 1 : 20-1 : 2000 (Flow cyt), (IL)  |
| <b>Expiration Date</b>     | 12 months from date of receipt.   |