
Product Datasheet

TRAcP Antibody / TRAP / ACP5 (orb749748)

Description

It recognizes a protein of 35kDa, which is identified as


Species/Host

Mouse

Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Tested Applications

IHC-P

Immunogen

Recombinant full-length human protein was used as the immunogen for the TRAcP antibody.

Preservatives

0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin (US sourced) and 0.05% sodium azide

Storage

Store the TRAcP antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

Note

For research use only

Application notes

Optimal dilution of the TRAcP antibody should be determined by the researcher. 1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.

Formula

0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide

Isotype

Mouse IgG2b, kappa

Clonality

Monoclonal

Purity

Protein G affinity chromatography

Uniprot ID

P13686

Hazard Information


This TRAcP antibody is available for research use only.

Dilution Range


Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT

Expiration Date

12 months from date of receipt.



IHC:
Formalin-fixed,
paraffin-embedded
h...



SDS-PAGE
Analysis of
Purified,
BSA-Free ...