

---

## Product Datasheet

### Transglutaminase 2 Antibody (orb749813)

**Description**

Recognizes a 77-85kDa protein, identified as cellular or


**Species/Host**

Mouse

**Reactivity**


Human

**Conjugation**

Unconjugated

**Tested Applications**

IHC-P



IHC:  
Formalin-fixed,  
paraffin-embedded  
h...

**Immunogen**


Recombinant full-length human protein was used as the immunogen for the Transglutaminase 2 antibody.

**Preservatives**

0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin (US sourced) and 0.05% sodium azide

**Storage**

Store the Transglutaminase 2 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).



IHC:  
Formalin-fixed,  
paraffin-embedded  
h...

**Note**

For research use only

**Application notes**

Optimal dilution of the Transglutaminase 2 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

**Formula**

0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide

**Isotype**

Mouse IgG1, kappa

**Clonality**

Monoclonal

**Purity**

Protein G affinity chromatography

**Uniprot ID**


**P21980**

**Hazard Information**

This Transglutaminase 2 antibody is available for research use only.

**Dilution Range**

Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT (1),Prediluted IHC only format: incubate for 30 min at RT (2)



IHC:  
Formalin-fixed,  
paraffin-embedded  
h...

**Expiration Date**

12 months from date of receipt.