

Product Datasheet

Hsp70/HSPA1A/HSPA1B Antibody (orb76326)



www.biorbyt.com

Description hts. Hsp70/HSPA1A/HSPA1B Antibody

Species/Host Rabbit

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested ICC, IHC, WB

Applications

Immunogen A synthetic peptide corresponding to a sequence

at the C-terminus of human Hsp70 (577-596aa VISWLDANTLAEKDEFEHKR), different from the related mouse sequence by four amino acids and from the related rat sequence by three amino

acids.

Form/Appearance Lyophilized

Concentration Adding 0.2 ml of distilled water will yield a

concentration of 500 µg/ml.

Storage Store at -20°C for one year from date of receipt.

After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Note For research use only

Application notes Tested Species: In-house tested species with

positive results. Predicted Species: Species predicted to be fit for the product based on sequence similarities. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins in acquiring of forms line and forms.

is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users. . Add 0.2ml of distilled water will yield a

concentration of 500ug/ml.

Isotype Rabbit IgG

Clonality Polyclonal

MW 70052 MW


Uniprot ID PODMV8

Dilution Range Immunohistochemistry (Paraffin-embedded

Section), 0.5-1µg/ml, Human, By Heat

Immunocytochemistry , 0.5-1µg/ml, Human, - Immunohistochemistry (Frozen Section), 0.5-1µg/ml, Rat, HumanWestern blot, 0.1-0.5µg/ml,


Human



Western blot analysis of Lane 1: 293T Ce...



Immunohistochemical analysis of formalin...



Carolina < br > 27709. United States

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North